

Satyridae16. *Maniola jurtina* Linnaeus

Generally distributed and common. Large specimens; males most often with orange-brown markings on FW upperside. Females near form *fortunata* Alpheraky, brightly coloured and with much extended orange-brown markings above.

17. *Lasiommata megera* Linnaeus

A single specimen recorded from near Parikia.

Hesperiidae18. *Carcharodus alceae* Esper

In fair numbers near Lefkes and near Aghios Minas.

19. *Thymelicus acteon* Rottemburg

A single male captured in Voutakou, at sea level.

20. *Gegenes pumilio* Hoffmannsegg

A few males captured on the stony bottom of a ravine, near Lefkes. Identification confirmed by the genitalia.

During a second visit to the island of Siphnos in April 1979 the following two species were recorded that had not been included in my previous list of 1978: *Papilio machaon* and *Gegenes pumilio*, both captured near Chrysopighi.

References

- Coutsis, J. G. 1978. Spring butterflies on the Greek island of Siphnos. *Entomologists's Rec.* **90**: 300-301.
Higgins, L. G. and Riley, N. D. 1980. *A Field Guide to the Butterflies of Britain and Europe*. Collins, London.
Polunin, O. 1969. *Flowers of Europe*. Oxford University Press, London.
Polunin, O. 1980. *Flowers of Greece and the Balkans*. Oxford University Press, Oxford.
-

THE PALE FORM OF THE PEARL BORDERED FRITILLARY: *BOLORIA EUPHROSYNE* L. AB. *PALLIDA* SPULER.—During a short stroll through Parkhurst Forest, Isle of Wight, on May 27 1981, I came upon a clearing where *B. euphrosyne* was flying plentifully, some of which were in fresh condition. After a few minutes I noticed a very pale example which immediately flew away rapidly. But it was then getting late so I resolved to return the following day. The next morning, having located the clearing, by a curious coincidence the first *euphrosyne* noted was the pale form and after netting it I saw it was a male ab. *pallida*.—F. H. CLOUTER, Helice, Glendale Road, Minster-in-Sheppey, Kent.